

Information video

An information video “Homeward Bound – Following Gynaecological Surgery” has also been produced which you can find on the National Women’s website at the following link:

<http://nationalwomenshealth.adhb.govt.nz/health-information/information-videos#Homeward>

Problems after discharge

Please contact your family doctor (GP) if you have any concerns. They will contact the hospital if necessary.

Resources

The Royal Australia & New Zealand College of Obstetricians & Gynaecologist online Patient Information: www.ranzcog.edu.au/womens-health/online-patient-information.html

Continence NZ: Healthy Bowel, Pelvic Floor exercise and other information.
www.continence.org.nz/pages/Continence-Information-Adults/18/

Te Whatu Ora
Health New Zealand
Te Toka Tumai Auckland

Classification number: WH135 (Reviewed July 24)
National Women’s Information Unit hiws@adhb.govt.nz

Te Whatu Ora
Health New Zealand

**Take Home
Advice Following
Gynaecological
Surgery**

Take regular pain relief

Panadol, Ibuprofen (or an alternative anti-inflammatory) and Tramadol are the most common medications prescribed on discharge from hospital.

Not everyone will be completely pain-free at discharge, so it is important that these medications are taken regularly for at least two weeks following your discharge from the hospital.

We generally recommend taking your anti-inflammatory medication as prescribed with food, and Panadol 3 to 4 times a day for 2 weeks unless advised differently.

In hospital you were given:

Paracetamol at _____

Anti-inflammatory at _____

Other Medication _____

Avoid constipation

Constipation is usually the main cause for discomfort and pain after surgery. You can avoid constipation by eating a diet of fruit (i.e. prunes, kiwifruit), vegetables, fibre, and lots of water after your surgery..

If you don't pass a bowel motion for 2 days after discharge from hospital, please get a laxative from your local pharmacy.

Take care of your wound

Keep your wound/wounds clean, and dry well after showering.

If your wound becomes red, painful, or starts oozing, you may be developing an infection, please see your GP.

Laparoscopic wounds

Take dressing off 2 days after surgery. If you have steristrips, leave them to fall off.

See your GP practice's nurse to remove stitches 7 days after surgery if they have not dissolved.

Laparoscopic hysterectomies

On discharge you will be given a surgical sponge. Shower with sponge after your dressings are removed.

Laparotomy wounds

Leave the dressing on until staples are removed 10 days after surgery. If dressing gets wet take it off and keep the wound dry. If there is any oozing from the wound or concerns please see your GP.

Staples are taken out 7-10 days after surgery. See your GP practice nurse to get staples removed. Your hospital nurse will give you a staple remover.

On discharge you will be given a surgical sponge. Shower with sponge after your dressings are removed.

Any other operation site dressings can be removed 3 days after surgery.

Vaginal bleeding (for hysterectomy patients)

You can expect some vaginal bleeding for 1-2 weeks after your operation. This is like a light period, and the discharge can be red or brown.

Many women have no or little bleeding straight after surgery and have a sudden gush of old blood or fluid approximately 10 days later. This usually stops quickly. Use sanitary pads, not tampons, as tampons increase the risk of infection.

Returning to activities

Depending on what procedure you have had, there will be a need to avoid unnecessary physical exertion in order to fully heal. Your surgeon will usually give specific advice but in general, if you have had an open operation with an abdominal scar, it will take six weeks before you can return to full activities.

A laparoscopic procedure has a shorter recovery time, usually two weeks.

Vaginal hysterectomies and procedures to repair prolapse will need a six week recovery, while tape operations for incontinence are faster, often being ready for normal activity at two weeks. Please ask your hospital doctor for specific advice before you leave hospital.